

A class of nonassociative algebras and cogebras

Michel Goze*

Laboratoire de Mathématiques et Applications,
Université de Haute Alsace,
4, rue des Frères Lumière, F-68093 Mulhouse Cedex, France.

Keywords: Nonassociative algebras, Lie-admissible algebras, Nonassociative cogebras

MS Classification numbers: 16Axx

Abstract

We present classes of nonassociative algebras whose associator satisfies invariance conditions given by the action of the 3 order symmetric group. Amongst these algebras we find the wellknown Vinberg algebras, the Pre-Lie algebras, the Lie-admissible algebras and the 3-power associative algebras.

I. Σ_3 -ASSOCIATIVE ALGEBRAS

A. Σ_3 -invariant spaces

Let Σ_3 be the 3-order symmetric group and $\mathbb{K}[\Sigma_3]$ the associated group algebra, where $\mathbb{K}$ is a field of characteristic zero. We denote by τ_{ij} be the transposition exchanging i and j , $c_1 = (1, 2, 3)$ and $c_2 = c_1^2$ the two 3-cycles of Σ_3 . Every $v \in \mathbb{K}[\Sigma_3]$ decomposes as follows:

$$v = a_1 id + a_2 \tau_{12} + a_3 \tau_{13} + a_4 \tau_{23} + a_5 c_1 + a_6 c_2$$

or simply

$$v = \sum_{\sigma \in \Sigma_3} a_\sigma \sigma$$

where $a_\sigma \in \mathbb{K}$.

Consider the natural right action of Σ_3 on $\mathbb{K}[\Sigma_3]$

$$\begin{aligned} \Sigma_3 \times \mathbb{K}[\Sigma_3] &\rightarrow \mathbb{K}[\Sigma_3] \\ (\sigma, \sum_i a_i \sigma_i) &\mapsto \sum_i a_i \sigma^{-1} \circ \sigma_i \end{aligned}$$

For every vector $v \in \mathbb{K}[\Sigma_3]$ we denote by $\mathcal{O}(v)$ the corresponding orbit of v . Let $F_v = \mathbb{K}(\mathcal{O}(v))$ be the linear subspace of $\mathbb{K}[\Sigma_3]$ generated by $\mathcal{O}(v)$. It is an invariant subspace of $\mathbb{K}[\Sigma_3]$. Therefore, using Mashke's theorem, we deduce that it is a direct sum of irreducible invariant subspaces.

B. Σ_3 -associative algebras

Let $(\mathcal{A}, \mu)$ be a $\mathbb{K}$ -algebra with multiplication μ . We denote by A_μ the associator of μ that is

$$A_\mu = \mu \circ (\mu \otimes Id - Id \otimes \mu).$$

*Email address: M.Goze@uha.fr

Every $\sigma \in \Sigma_3$ defines a linear map denoted by Φ_σ given by

$$\begin{array}{ccc} \Phi_\sigma : & \mathcal{A}^{\otimes 3} & \rightarrow \\ & x_1 \otimes x_2 \otimes x_3 & \rightarrow x_{\sigma^{-1}(1)} \otimes x_{\sigma^{-1}(2)} \otimes x_{\sigma^{-1}(3)}. \end{array}$$

If $v = \sum_{\sigma \in \Sigma_3} a_\sigma \sigma$ is a vector of $\mathbb{K}[\Sigma_3]$ we define the endomorphism Φ_v of $\mathcal{A}^{\otimes 3}$ by taking:

$$\Phi_v = \sum_{\sigma \in \Sigma_3} a_\sigma \Phi_\sigma.$$

Definition 1 *An algebra $(\mathcal{A}, \mu)$ is a Σ_3 -associative algebra if there exists $v \in \mathbb{K}[\Sigma_3]$ such that*

$$A_\mu \circ \Phi_v = 0.$$

C. Lie-admissible algebras. 3-power associative algebras

Proposition 2 *Let v be in $\mathbb{K}[\Sigma_3]$ such that $\dim F_v = 1$. Then $v = \alpha V$ or $v = \alpha W$ with $\alpha \in \mathbb{K}$ where the vectors V and W are the following vectors :*

$$V = Id - \tau_{12} - \tau_{13} - \tau_{23} + c_1 + c_2, \quad (1)$$

$$W = Id + \tau_{12} + \tau_{13} + \tau_{23} + c_1 + c_2. \quad (2)$$

The first case corresponds to the character of Σ_3 given by the signature, the second corresponds to the trivial case.

Every algebra $(\mathcal{A}, \mu)$ whose associator satisfies

$$A_\mu \circ \Phi_V = 0$$

is a Lie-admissible algebra. This means that the algebra $(\mathcal{A}, [,])$ whose product is given by the bracket $[x, y] = \mu(x, y) - \mu(y, x)$ is a Lie algebra. Likewise an algebra $(\mathcal{A}, \mu)$ whose associator satisfies

$$A_\mu \circ \Phi_W = 0$$

is 3-power associative that is it satisfies $A_\mu(x, x, x) = 0$ for every $x \in A$.

II. G_i -ASSOCIATIVE ALGEBRAS

In this section we study the Σ_3 -associative algebras corresponding to the subgroups of Σ_3 .

A. Notations

Let us consider the subgroups of Σ_3 :

$$\begin{aligned} G_1 &= \{Id\}, \\ G_2 &= \{Id, \tau_{12}\}, \\ G_3 &= \{Id, \tau_{23}\}, \\ G_4 &= \{Id, \tau_{13}\}, \\ G_5 &= \{Id, c_1, c_2\} = A_3 \text{ (the alternating group)}, \\ G_6 &= \Sigma_3 \end{aligned}$$

We keep these notations for the subgroups of Σ_3 in the following sections.

B. G_i -associative algebras

This notion was introduced by E.Remm in her thesis [10].

Definition 3 *Let G_i be a subgroup of Σ_3 . The algebra $(\mathcal{A}, \mu)$ is G_i -associative if*

$$\sum_{\sigma \in G_i} (-1)^{\varepsilon(\sigma)} A_\mu \circ \Phi_\sigma = 0.$$

Proposition 4 *Every G_i -associative algebra is a Σ_3 -associative algebra.*

Proof. Every subgroup G_i of Σ_3 corresponds to an invariant linear space $F(v_i)$ generated by a single vector $v_i \in \mathbb{K}[\Sigma_3]$. More precisely we consider $v_1 = Id$, $v_2 = \tau_{12}$, $v_3 = \tau_{23}$, $v_4 = \tau_{13}$, $v_5 = Id + c_1 + c_2$ and $v_6 = V$ that we have defined in (1). $\square$

Proposition 5 *Every G_i -associative algebra is a Lie-admissible algebra.*

Proof. The vector V belongs to the orbits $\mathcal{O}(v_i)$ for every v_i . Thus, if μ is a G_i -associative product, it also satisfies

$$A_\mu \circ \Phi_V = 0$$

and μ is a Lie-admissible multiplication. $\square$

We deduce the following type of Lie-admissible algebras :

1. A G_1 -associative algebra is an associative algebra.
2. A G_2 -associative algebra is a Vinberg algebra. If A is finite-dimensional, the associated Lie algebra is provided with an affine structure.
3. A G_3 -associative algebra is a pre-Lie algebra.
4. If (A, μ) is G_4 -associative then μ satisfies

$$(X.Y).Z - X.(Y.Z) = (Z.Y).X - Z.(Y.X)$$

with $X.Y = \mu(X, Y)$.

5. If (A, μ) is G_5 -associative then μ satisfies the generalized Jacobi condition :

$$(X.Y).Z + (Y.Z).X + (Z.X).Y = X.(Y.Z) + Y.(Z.X) + Z.(X.Y)$$

with $X.Y = \mu(X, Y)$. Moreover if the product is antisymmetric, then it is a Lie algebra bracket.

6. A G_6 -associative algebra is a Lie-admissible algebra.

III. G_i -ASSOCIATIVE COGEBRAS

A. An axiomatic definition of the G_i -associative algebras

To define a G_i -cogebra it is easier to present an equivalent and axiomatic definition of the notion of G_i -associative algebra.

Definition 6 *A G_i -associative algebra is (A, μ, η, G_i) where A is a vector space, G_i a subgroup of Σ_3 , $\mu : A \otimes A \longrightarrow A$ and $\eta : \mathbb{K} \rightarrow A$ are linear maps satisfying the following axioms :*

1. (G_i -ass): The square

$$\begin{array}{ccc} A \otimes A \otimes A & \xrightarrow{(\mu \otimes Id)_{G_i}} & A \otimes A \\ (Id \otimes \mu)_{G_i} \downarrow & & \downarrow \mu \\ A \otimes A & \xrightarrow{\mu} & A \end{array}$$

commutes, where $(Id \otimes \mu)_{G_i}$ is the linear mapping defined by:

$$(Id \otimes \mu)_{G_i} = \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} (Id \otimes \mu) \circ \Phi_\sigma.$$

2. (Un) The following diagram is commutative :

$$\begin{array}{ccccc} \mathbb{K} \otimes A & \xrightarrow{\eta \otimes id} & A \otimes A & \xleftarrow{id \otimes \eta} & A \otimes \mathbb{K} \\ & \searrow & \downarrow \mu & \swarrow & \\ & & A & & \end{array}$$

The axiom (G_i -ass) expresses that the multiplication μ is G_i -associative whereas the axiom (Un) means that the element $\eta(1)$ of $\mathcal{A}$ is a left and right unit for μ .

Definition 7 A morphism of G_i -associative algebras

$$f : (\mathcal{A}, \mu, \eta, G_i) \rightarrow (\mathcal{A}', \mu', \eta', G_i)$$

is a linear map from $\mathcal{A}$ to $\mathcal{A}'$ such that

$$\mu' \circ (f \otimes f) = f \circ \mu \quad \text{and} \quad f \circ \eta = \eta'$$

B. G_i -associative cogebras

We want to dualize the previous diagrams to obtain the notions of corresponding cogebras. Let Δ be a comultiplication on a vector space C :

$$\Delta : C \longrightarrow C \otimes C.$$

We define the bilinear map

$$G_i \circ (\Delta \otimes Id) : C^{\otimes 3} \longrightarrow C^{\otimes 3}$$

by

$$G_i \circ (\Delta \otimes Id) = \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} \Phi_\sigma \circ (\Delta \otimes Id).$$

Definition 8 A G_i -cogebra is a vector space C provided with a comultiplication $\Delta : C \longrightarrow C \otimes C$ and a counit $\epsilon : C \rightarrow \mathbb{K}$ such that :

1. (G_i -ass co) The following square is commutative :

$$\begin{array}{ccc} C & \xrightarrow{\Delta} & C \otimes C \\ \Delta \downarrow & & \downarrow G_i \circ (Id \otimes \Delta) \\ C \otimes C & \xrightarrow{G_i \circ (\Delta \otimes Id)} & C \otimes C \otimes C. \end{array}$$

2. (Coun) The following diagram is commutative :

$$\begin{array}{ccccc} & & \mathbb{K} \otimes C & \xleftarrow{\epsilon \otimes id} & C \otimes C & \xrightarrow{id \otimes \epsilon} & C \otimes \mathbb{K} \\ & & \swarrow & & \uparrow \Delta & & \searrow \\ & & C & & C & & \end{array}$$

Definition 9 A morphism of G_i -associative cogebras

$$f : (C, \Delta, \epsilon, G_i) \rightarrow (C', \Delta', \epsilon', G_i)$$

is a linear map from C to C' such that

$$(f \otimes f) \circ \Delta = \Delta' \circ f \quad \text{and} \quad \epsilon = \epsilon' \circ f$$

C. The dual space of a G_i -associative cogebra

For any natural number n and any $\mathbb{K}$ -vector spaces E and F , we denote by

$$\lambda_n : Hom(E, F)^{\otimes n} \longrightarrow Hom(E^{\otimes n}, F^{\otimes n})$$

the natural embedding

$$\lambda_n(f_1 \otimes \dots \otimes f_n)(x_1 \otimes \dots \otimes x_n) = f_1(x_1) \otimes \dots \otimes f_n(x_n).$$

Proposition 10 The dual space of a G_i -associative cogebra is a G_i -associative algebra.

Proof. Let (C, Δ) a G_i -associative cogeбра. We consider the multiplication on the dual vector space C^* of C defined by :

$$\mu = \Delta^* \circ \lambda_2.$$

It provides C^* with a G_i -associative algebra structure. In fact we have

$$\mu(f_1 \otimes f_2) = \mu_{\mathbb{K}} \circ \lambda_2(f_1 \otimes f_2) \circ \Delta \quad (3)$$

for all $f_1, f_2 \in C^*$ where $\mu_{\mathbb{K}}$ is the multiplication of $\mathbb{K}$. The equation (3) becomes :

$$\begin{aligned} \mu \circ (\mu \otimes Id)(f_1 \otimes f_2 \otimes f_3) &= \mu_{\mathbb{K}} \circ (\lambda_2(\mu(f_1 \otimes f_2) \otimes f_3) \circ \Delta \\ &= \mu_{\mathbb{K}} \circ \lambda_2((\mu_{\mathbb{K}} \circ \lambda_2(f_1 \otimes f_2) \circ \Delta) \otimes f_3) \circ \Delta \\ &= \mu_{\mathbb{K}} \circ (\mu_{\mathbb{K}} \otimes Id) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (\Delta \otimes Id) \circ \Delta. \end{aligned}$$

The associator A_{μ} satisfies :

$$\begin{aligned} A_{\mu} &= \mu_{\mathbb{K}} \circ (\mu_{\mathbb{K}} \otimes Id) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (\Delta \otimes Id) \circ \Delta \\ &\quad - \mu_{\mathbb{K}} \circ (Id \otimes \mu_{\mathbb{K}}) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (Id \otimes \Delta) \circ \Delta. \end{aligned}$$

and using associativity and commutativity of the multiplication in $\mathbb{K}$, we obtain

$$A_{\mu} = \mu_{\mathbb{K}} \circ (\mu_{\mathbb{K}} \otimes Id) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ ((\Delta \otimes Id) \circ \Delta - (Id \otimes \Delta) \circ \Delta).$$

For every $\sigma \in \Sigma_3$ and any $\mathbb{K}$ -vector space E , we denote by Φ_{σ}^E the action

$$\Phi_{\sigma}^E : E^{\otimes 3} \longrightarrow E^{\otimes 3}$$

defined in the first section :

$$\Phi_{\sigma}^E(e_1 \otimes e_2 \otimes e_3) = e_{\sigma^{-1}(1)} \otimes e_{\sigma^{-1}(2)} \otimes e_{\sigma^{-1}(3)}.$$

Thus

$$\begin{aligned} &\sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_{\mu} \circ \Phi_{\sigma}^{C^*} \\ &= \mu_{\mathbb{K}} \circ (\mu_{\mathbb{K}} \otimes Id) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (G_i \circ (\Delta \otimes Id) \circ \Delta - G_i \circ (Id \otimes \Delta) \circ \Delta) \\ &= 0. \end{aligned}$$

□

D. The dual space of a finite dimensional G_i -associative algebra

Proposition 11 *The dual vector space of a finite dimensional G_i -associative algebra has a G_i -associative cogeбра structure.*

Proof. Let $\mathcal{A}$ be a finite dimensional G_i -associative algebra and let $\{e_i, i = 1, \dots, n\}$ be a basis of $\mathcal{A}$. If $\{f_i\}$ is the dual basis then $\{f_i \otimes f_j\}$ is a basis of $\mathcal{A}^* \otimes \mathcal{A}^*$. The coproduct Δ on $\mathcal{A}^*$ is defined by

$$\Delta(f) = \sum_{i,j} f(\mu(e_i \otimes e_j)) f_i \otimes f_j.$$

In particular

$$\Delta(f_k) = \sum_{i,j} C_{ij}^k f_i \otimes f_j$$

where C_{ij}^k are the structure constants of μ related to the basis $\{e_i\}$. Then Δ is the comultiplication of a G_i -associative cogebras.

E. Associated Lie cogebras

Since every G_i -associative algebra $(\mathcal{A}, \mu)$ is Lie-admissible, the bilinear map $\mu(X, Y) - \mu(Y, X)$ is a Lie bracket. Similarly we prove a similar property for G_i -associative cogebras. Let us recall the notion of Lie cogebras. Let C un $\mathbb{K}$ -vector space and

$$\Delta : C \longrightarrow C \otimes C$$

a linear map satisfying

1. $\tau \circ \Delta = -\Delta$ where τ is the permutation $\tau(x \otimes y) = y \otimes x$
2. $\Phi_v \circ (Id \otimes \Delta) \circ \Delta = 0$ where $v = \sum_{\sigma \in G_5} \sigma$.

Then (C, Δ) is called a Lie cogebras.

Proposition 12 *Let (C, Δ) be a G_i -associative cogebras (non necessary counitary). Let Δ_L the linear map defined by $\Delta_L = \Delta - \tau \circ \Delta$. Then (C, Δ_L) is a Lie cogebras.*

Proof. It is clear that $\tau \circ \Delta_L = -\Delta_L$. As W is in the orbit of the vector $u_i = \sum_{\sigma \in G_i} \sigma$, then any G_i -associative cogebras is G_6 -associative cogebras, that is a Lie admissible cogebras. But the identity

$$\Phi_W \circ (Id \otimes \Delta) \circ \Delta - \Phi_W \circ (\Delta \otimes Id) \circ \Delta = 0$$

is equivalent to

$$\begin{aligned} & \Phi_v \circ (Id \otimes \Delta) \circ \Delta - \tau \circ \Phi_v \circ (Id \otimes \Delta) \circ \Delta \\ & - \Phi_v \circ (Id \otimes \tau \circ \Delta) \circ \Delta - \tau \circ \Phi_v \circ (Id \otimes \tau \circ \Delta) \circ \Delta = 0 \end{aligned}$$

with $v = Id + c + c^2$, we deduce the result. $\square$

IV. THE CONVOLUTION PRODUCT

Let $(\mathcal{A}, \mu, \eta, G_1)$ be an associative algebra and $(C, \Delta, \epsilon, G_1)$ an associative cogebras. The convolution product on the vector space $Hom(C, \mathcal{A})$ is given by

$$f * g = \mu \circ \lambda_2 \circ (f \otimes g) \circ \Delta$$

for every f and $g \in Hom(C, \mathcal{A})$ and $(Hom(C, \mathcal{A}), \star, G_1)$ is an associative algebra. We want to generalize this property to the other groups G_i , $i \geq 2$. Let us begin by introduce new classes of associative algebras which appear naturally by studying the duality in the corresponding operads (see section V).

A. The $G_i^!$ -algebras and cogebras

Definition 13 *For $i \geq 2$, a $G_i^!$ -algebra is an associative algebra satisfying :*

- for $i = 2$: $x_1.x_2.x_3 = x_2.x_1.x_3$
- for $i = 3$: $x_1.x_2.x_3 = x_1.x_3.x_2$
- for $i = 4$: $x_1.x_2.x_3 = x_3.x_2.x_1$
- for $i = 5$: $x_1.x_2.x_3 = x_2.x_3.x_1 = x_3.x_1.x_2$
- for $i = 6$: $x_1.x_2.x_3 = x_{\sigma(1)}.x_{\sigma(2)}.x_{\sigma(3)}$ for all x_1, x_2, x_3 and $\sigma \in \Sigma_3$.

Definition 14 *For $i \geq 2$, a $G_i^!$ -cogebras is a coassociative cogebras, that is a G_1 -associative cogebras, satisfying :*

$$\Phi_{u_i} \circ (Id \otimes \Delta) \circ \Delta = (Id \otimes \Delta) \circ \Delta$$

where $u_i = \sum_{\sigma \in G_i} \sigma^{-1}$.

B. Convolution product on $\text{Hom}(C, \mathcal{A})$

We will provide $\text{Hom}(C, \mathcal{A})$ with a structure of G_i -associative algebra.

Proposition 15 *Let $(\mathcal{A}, \mu, \eta, G_i)$ be a G_i -associative algebra and $(C, \Delta, \epsilon, G_i^!)$ a $G_i^!$ -cogebra. Then $(\text{Hom}(C, \mathcal{A}), \star, \eta \circ \epsilon, G_i)$ is a G_i -associative algebra where $\star$ is the convolution product :*

$$f \star g = \mu \circ \lambda_2(f \otimes g) \circ \Delta.$$

Proof. Let us compute the associator $A_\star$ of the convolution product.

$$\begin{aligned} (f_1 \star f_2) \star f_3 &= \mu \circ \lambda_2((f_1 \star f_2) \otimes f_3) \circ \Delta \\ &= \mu \circ \lambda_2((\mu \circ \lambda_2(f_1 \otimes f_2) \circ \Delta) \otimes f_3) \circ \Delta \\ &= \mu \circ (\mu \otimes \text{Id}) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (\Delta \otimes \text{Id}) \circ \Delta. \end{aligned}$$

Thus

$$\begin{aligned} A_\star(f_1 \otimes f_2 \otimes f_3) &= \mu \circ (\mu \otimes \text{Id}) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (\Delta \otimes \text{Id}) \circ \Delta \\ &\quad - \mu \circ (\text{Id} \otimes \mu) \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (\text{Id} \otimes \Delta) \circ \Delta. \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_\star \circ \Phi_\sigma^{\text{Hom}(C, \mathcal{A})}(f_1 \otimes f_2 \otimes f_3) \\ = \mu \circ (\mu \otimes \text{Id}) \circ (\sum_{\sigma \in G_i} \lambda_3(\Phi_\sigma^{\text{Hom}(C, \mathcal{A})}(f_1 \otimes f_2 \otimes f_3))) \circ (\Delta \otimes \text{Id}) \circ \Delta \\ - \mu \circ (\text{Id} \otimes \mu) \circ (\sum_{\sigma \in G_i} \lambda_3(\Phi_\sigma^{\text{Hom}(C, \mathcal{A})}(f_1 \otimes f_2 \otimes f_3))) \circ (\text{Id} \otimes \Delta) \circ \Delta. \end{aligned}$$

But

$$\lambda_3(\Phi_\sigma^{\text{Hom}(C, \mathcal{A})}(f_1 \otimes f_2 \otimes f_3)) = \Phi_\sigma^A \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ \Phi_{\sigma^{-1}}^C.$$

This gives

$$\begin{aligned} \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_\star \circ \Phi_\sigma^{\text{Hom}(C, \mathcal{A})}(f_1 \otimes f_2 \otimes f_3) \\ = \mu \circ (\mu \otimes \text{Id}) \circ (\sum_{\sigma \in G_i} \Phi_\sigma^A \circ \lambda_3(f_1 \otimes f_2 \otimes f_3)) \circ \Phi_{\sigma^{-1}}^C \circ (\Delta \otimes \text{Id}) \circ \Delta \\ - \mu \circ (\text{Id} \otimes \mu) \circ (\sum_{\sigma \in G_i} \Phi_\sigma^A \circ \lambda_3(f_1 \otimes f_2 \otimes f_3)) \circ \Phi_{\sigma^{-1}}^C \circ (\text{Id} \otimes \Delta) \circ \Delta. \end{aligned}$$

As Δ is coassociative

$$(\Delta \otimes \text{Id}) \circ \Delta = (\text{Id} \otimes \Delta) \circ \Delta$$

and the $G_i^!$ -cogebra structure implies

$$\Phi_{u_i}^C \circ (\text{Id} \otimes \Delta) \circ \Delta = \Phi_{u_i}^C \circ (\Delta \otimes \text{Id}) \circ \Delta = (\Delta \otimes \text{Id}) \circ \Delta.$$

Then

$$\begin{aligned} \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_\star \circ \Phi_\sigma^{\text{Hom}(C, \mathcal{A})}(f_1 \otimes f_2 \otimes f_3) \\ = \mu \circ (\mu \otimes \text{Id}) \circ (\sum_{\sigma \in G_i} \Phi_\sigma^A \circ \lambda_3(f_1 \otimes f_2 \otimes f_3)) \circ (\Delta \otimes \text{Id}) \circ \Delta \\ - \mu \circ (\text{Id} \otimes \mu) \circ (\sum_{\sigma \in G_i} \Phi_\sigma^A \circ \lambda_3(f_1 \otimes f_2 \otimes f_3)) \circ (\Delta \otimes \text{Id}) \circ \Delta \\ = \sum_{\sigma \in G_i} A_\mu \circ \Phi_\sigma^A \circ \lambda_3(f_1 \otimes f_2 \otimes f_3) \circ (\Delta \otimes \text{Id}) \circ \Delta \\ = 0. \end{aligned}$$

This proves the proposition. $\square$

V. TENSOR PRODUCT of Σ_3 -ASSOCIATIVE ALGEBRAS

A. Non monoidal categories We know that the tensor product of associative algebras can be provided with an associative algebra structure. In other words, the category of associative algebras is monoidal and closed for the tensor product. This is generally not true for other categories of Σ_3 -associative algebras.

Proposition 16 *Let $(\mathcal{A}, \mu_A)$ and $(\mathcal{B}, \mu_B)$ be two Σ_3 -associative algebras respectively defined by the relations $A_{\mu_A} \circ \Phi_v = 0$ and $A_{\mu_B} \circ \Phi_w = 0$. Then $(\mathcal{A} \otimes_{\mathbb{K}} \mathcal{B}, \mu_A \otimes \mu_B)$ is a Σ_3 -associative algebra if and only if $\mathcal{A}$ and $\mathcal{B}$ are associative algebras (i.e G_1 -associative algebras).*

Proof. Let A_μ be the associator of the law μ and $A_\mu^L(x_1 \otimes x_2 \otimes x_3) = \mu(\mu(x_1, x_2), x_3)$ and $A_\mu^R = A_\mu^L - A_\mu$. By hypothesis, $\mathcal{A}$ (resp. $\mathcal{B}$) is defined by $A_{\mu_A} \circ \Phi_v = 0$ (resp. $A_{\mu_B} \circ \Phi_w = 0$). Let us suppose that $\mathcal{A} \otimes \mathcal{B}$ is a Σ_3 -associative algebra. There exists $v' \in \mathbb{K}$ such that $A_{\mu_{\mathcal{A} \otimes \mathcal{B}}} \circ \Phi_{v'} = 0$. By taking $v' = \sum_{i=1}^6 \gamma_i \sigma_i$ the last condition becomes

$$\sum_{i=1}^6 \gamma_i [A_{\mu_{\mathcal{A} \otimes \mathcal{B}}} \circ \Phi_{\sigma_i}] = 0 \quad (3)$$

and

$$\sum_{i=1}^6 \gamma_i [A_{\mu_A}^L \circ \Phi_{\sigma_i} \otimes A_{\mu_B}^L \circ \Phi_{\sigma_i} - A_{\mu_A}^R \circ \Phi_{\sigma_i} \otimes A_{\mu_B}^R \circ \Phi_{\sigma_i}] = 0. \quad (4)$$

Let us denote by $e_i = A_{\mu_A}^L \circ \Phi_{\sigma_i}$, $\tilde{e}_i = A_{\mu_A}^R \circ \Phi_{\sigma_i}$, $f_i = A_{\mu_B}^L \circ \Phi_{\sigma_i}$ and $\tilde{f}_i = A_{\mu_B}^R \circ \Phi_{\sigma_i}$. The vectors e_i and $\tilde{e}_i$ belong to $Hom(\mathcal{A}^{\otimes 3}, \mathcal{A})$ and the vectors f_i and $\tilde{f}_i$ to $Hom(\mathcal{B}^{\otimes 3}, \mathcal{B})$. Equation 4 becomes :

$$\sum_{i=1}^6 \gamma_i [e_i \otimes f_i - \tilde{e}_i \otimes \tilde{f}_i] = 0. \quad (5)$$

From the definition of the algebras $\mathcal{A}$ and $\mathcal{B}$ we have:

$$\sum_{i=1}^6 a_i (e_i - \tilde{e}_i) = 0 \text{ and } \sum_{i=1}^6 b_i (f_i - \tilde{f}_i) = 0$$

if $v = \sum_{i=1}^6 a_i \sigma_i$ and $w = \sum_{i=1}^6 b_i \sigma_i$. Suppose that $\dim F_w = k$ with $0 \leq k \leq 6$. Then the rank of the vectors $\{f_i, \tilde{f}_i\}$ is equal to $(6 + k)$. We can suppose that $\{f_1, \dots, f_6, \tilde{f}_1, \dots, \tilde{f}_k\}$ are independent. This implies:

$$\begin{cases} \tilde{f}_{k+1} = \rho_1^{k+1} f_1 + \dots + \rho_6^{k+1} f_6 + \tilde{\rho}_1^{k+1} \tilde{f}_1 + \dots + \tilde{\rho}_k^{k+1} \tilde{f}_k \\ \vdots \\ \tilde{f}_6 = \rho_1^6 f_1 + \dots + \rho_6^6 f_6 + \tilde{\rho}_1^6 \tilde{f}_1 + \dots + \tilde{\rho}_k^6 \tilde{f}_k. \end{cases}$$

The equation 5 becomes:

$$\sum_{i=1}^6 e'_i \otimes f_i + \sum_{i=1}^k e''_i \otimes \tilde{f}_k = 0$$

and the independence of the vectors $\{f_1, \dots, f_6, \tilde{f}_1, \dots, \tilde{f}_k\}$ implies $e'_1 = \dots = e'_6 = e''_1 = \dots = e''_k = 0$. Then we deduce

$$\begin{cases} \gamma_1 e_1 - \gamma_{k+1} \rho_1^{k+1} \tilde{e}_1 - \dots - \gamma_6 \rho_1^6 \tilde{e}_6 = 0 \\ \vdots \\ \gamma_6 e_6 - \gamma_{k+1} \rho_6^{k+1} \tilde{e}_1 - \dots - \gamma_6 \rho_6^6 \tilde{e}_6 = 0 \\ \gamma_1 \tilde{e}_1 + \gamma_{k+1} \tilde{\rho}_1^{k+1} \tilde{e}_{k+1} + \dots + \gamma_6 \tilde{\rho}_1^6 \tilde{e}_6 = 0 \\ \vdots \\ \gamma_k \tilde{e}_k + \gamma_{k+1} \tilde{\rho}_k^{k+1} \tilde{e}_{k+1} + \dots + \gamma_6 \tilde{\rho}_k^6 \tilde{e}_6 = 0. \end{cases}$$

Let us suppose that $k \neq 0$. Then the conditions on the vector v are of the type $\sum_{i=1}^6 a_i(e_i - \tilde{e}_i) = 0$ and we have successively $\gamma_1 = \dots = \gamma_k = 0$ and $\gamma_{k+1} \tilde{\rho}_i^{k+1} = \dots = \gamma_6 \tilde{\rho}_i^6 = 0$ for $i = 1, \dots, 6$. If one of γ_j for $j = k+1, \dots, 6$ is not 0 then $\tilde{\rho}_j^i = 0$ for $i = 1, \dots, 6$. Thus

$$\tilde{f}_j = \rho_1^j f_1 + \dots + \rho_6^j f_6$$

and, as for every i there exists $\sigma \in \Sigma_3$ such that $\tilde{f}_i = \tilde{f}_j \circ \Phi_\sigma$, we have also

$$\tilde{f}_i = \rho_1^i f_1 + \dots + \rho_6^i f_6$$

for every $i = 1, \dots, 6$. This implies $k = 0$ which is impossible from the hypothesis. Thus $k = 0$ and the vectors $\{f_i, \tilde{f}_i\}_{i=1, \dots, 6}$ is of rank 6. The only possible relations are then $f_i = \tilde{f}_i$ and $\mathcal{B}$ is associative. We deduce the associativity of $\mathcal{A}$. $\square$

B. $G_i!$ -algebras

In [7], we have studied the operads corresponding to the G_i -associative algebras. More precisely, the operad $G_i - \mathcal{A}ss$ is the quadratic operad defined by the relations

$$\sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} (x_{\sigma^{-1}(1)} \cdot x_{\sigma^{-1}(2)}) \cdot x_{\sigma^{-1}(3)} - x_{\sigma^{-1}(1)} \cdot (x_{\sigma^{-1}(2)}) \cdot x_{\sigma^{-1}(3)}).$$

Then an algebra on the operad $G_i - \mathcal{A}ss$ is a G_i -associative algebra. In [7], we have also computed the dual operads $(G_i - \mathcal{A}ss)!$ and proved that only $G_1 - \mathcal{A}ss$ (classically noted $\mathcal{A}ss$ is autodual that is $(G_i - \mathcal{A}ss) = (G_i - \mathcal{A}ss)!$. We call a $G_i!$ -algebra an algebra on the quadratic operad $(G_i - \mathcal{A}ss)!$. In definition 13 we have given their structures.

C. Tensor products

Theorem 17 *If $\mathcal{A}$ is a G_i -associative algebra and $\mathcal{B}$ a $G_i!$ -algebra (with the same indice) then $\mathcal{A} \otimes \mathcal{B}$ can be provided with a G_i -algebra structure for $i = 1, \dots, 6$.*

Proof. Let us consider on $\mathcal{A} \otimes \mathcal{B}$ the classical tensor product

$$\mu_A \otimes \mu_B((a_1 \otimes b_1) \otimes (a_2 \otimes b_2)) = \mu_A(a_1 \otimes a_2) \otimes \mu_B(b_1 \otimes b_2).$$

To simplify, we denote by μ the product $\mu_A \otimes \mu_B$. As $\mathcal{B}$ is an associative algebra, the associator A_μ satisfies:

$$A_\mu((a_1 \otimes b_1) \otimes (a_2 \otimes b_2)) \otimes (a_3 \otimes b_3) = A_{\mu_A}(a_1 \otimes a_2 \otimes a_3) \otimes \mu_B \circ (\mu_B \otimes Id)(b_1 \otimes b_2 \otimes b_3).$$

Therefore

$$\begin{aligned} \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_\mu \circ \Phi_\sigma((a_1 \otimes b_1) \otimes (a_2 \otimes b_2) \otimes (a_3 \otimes b_3)) \\ = \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_{\mu_A} \circ \Phi_\sigma(a_1 \otimes a_2 \otimes a_3) \otimes \mu_B \circ (\mu_B \otimes Id) \circ \Phi_\sigma(b_1 \otimes b_2 \otimes b_3). \end{aligned}$$

But $\mathcal{B}$ a $G_i^!$ -algebra. Then

$$\mu_B \circ (\mu_B \otimes Id) \circ \Phi_\sigma(b_1 \otimes b_2 \otimes b_3) = \mu_B \circ (\mu_B \otimes Id)(b_1 \otimes b_2 \otimes b_3).$$

We obtain

$$\begin{aligned} & \sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_\mu \circ \Phi_\sigma((a_1 \otimes b_1) \otimes (a_2 \otimes b_2) \otimes (a_3 \otimes b_3)) \\ &= (\sum_{\sigma \in G_i} (-1)^{\epsilon(\sigma)} A_{\mu_A} \circ \Phi_\sigma(a_1 \otimes a_2 \otimes a_3)) \otimes \mu_B \circ (\mu_B \otimes Id)(b_1 \otimes b_2 \otimes b_3) \\ &= 0. \end{aligned}$$

□

D. The categories $G_i - ASS$

Let $G_i - ASS$ and $(G_i - ASS)^!$ the categories whose objects are respectively G_i -associative algebras and $G_i^!$ -algebras and the morphismes are the homomorphismes of algebras. The previous theorem can be interpreted as follows: Let $\mathcal{A}$ be a G_i -associative algebra. Then $\mathcal{A} \otimes -$ is a covariant functor

$$\mathcal{A} \otimes - : (G_i - ASS)^! \longrightarrow G_i - ASS.$$

Thanks: I thank the referee to point up a mistake on the section IV in the first version and to propose a correction.

References

- [1] Albert A.A., Power-associative rings. *Trans. Amer. Math. Soc.* **64** (1948), 552-593.
- [2] Bremmer M., Hentzel I., Identities for the associator in alternative algebras. Preprint.
- [3] Dzhumadil'daev A., Novikov-Jordan algebras. *Comm. in Algebra* **11**, (2002), 5207-5240.
- [4] Gerstenhaber M., The cohomology structure of an associative ring, *Ann of math.* **78**, 2, (1963), 267-288.
- [5] Ginzburg V., Kapranov M., Koszul duality for operads. *Duke Math Journal.* **76** 1, (1994), 203-272.
- [6] Elduque A., Myung H.C., *Mutations of Alternative Algebras*. Kluwer Academic Press. Math and its applications 278. 1994.
- [7] Goze M., Remm E., Lie-admissible algebras and operads. *Journal of Algebra* **273/1** (2004), 129-152.
- [8] Goze M., Remm E., A class of nonassociative algebras. Preprint Mulhouse 2003 and xxx RA/0309015
- [9] Nijenhuis A., Sur une classe de propriétés communes à quelques types différents d'algèbres. *Enseignement Math.* **14** (2), (1970), 225-277.
- [10] Remm E., Opérades Lie-admissibles. *C. R. Math. Acad. Sci. Paris* **334** no. 12, (2002), 1047-1050.
- [11] Schafer R.D. *Introduction to Nonassociative Algebras*. Academic Press. 1966